

**AMENDMENTS TO THE SPECIFICATION:**

Please amend the sixth paragraph on page 4 as follows:

D<sup>1</sup> --As indicated above, these ring systems can be unsubstituted or substituted by substituents such as halogen up to per-halosubstitution. Other suitable substituents for the moieties of B include alkyl, alkoxy, carboxy, cycloalkyl, aryl, heteroaryl, cyano, hydroxy and amine. These other substituents, generally referred to as X and X' herein, include -CN, -CO<sub>2</sub>R<sup>5</sup>, -C(O)NR<sup>5</sup>R<sup>5'</sup>, -C(O)R<sup>5</sup>, -NO<sub>2</sub>, -OR<sup>5</sup>, -SR<sup>5</sup>, -NR<sup>5</sup>R<sup>5'</sup>, -NR<sup>5</sup>C(O)OR<sup>5'</sup>, -NR<sup>5</sup>C(O)R<sup>5'</sup>, C<sub>1</sub>-C<sub>10</sub> alkyl, C<sub>2-10</sub>-alkenyl, C<sub>1-10</sub>-alkoxy, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>6</sub>-C<sub>14</sub> aryl, C<sub>7</sub>-C<sub>24</sub> alkaryl, C<sub>3</sub>-C<sub>13</sub> heteroaryl, C<sub>4</sub>-C<sub>23</sub> alkheteroaryl, substituted C<sub>1</sub>-C<sub>10</sub> alkyl, substituted C<sub>2-10</sub>-alkenyl, substituted C<sub>2-10</sub>-alkoxy, substituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl, substituted C<sub>4</sub>-C<sub>23</sub> alkheteroaryl and ~~Y AR~~ M-L<sup>1</sup>..--

Please amend the third and fourth paragraphs on page 5 as follows:

D<sup>2</sup> --The bridging group ~~Y M~~ is preferably -O-, -S-, -N(R<sup>5</sup>)-, -(CH<sub>2</sub>)<sub>m</sub>-, -C(O)-, -CH(OH)-, -NR<sup>5</sup>C(O)NR<sup>5</sup>R<sup>5'</sup>-, -NR<sup>5</sup>C(O)-, -C(O)NR<sup>5</sup>-, -(CH<sub>2</sub>)<sub>m</sub>O-, -(CH<sub>2</sub>)<sub>m</sub>S-, -(CH<sub>2</sub>)<sub>m</sub>N(R<sup>5</sup>)-, -O(CH<sub>2</sub>)<sub>m</sub>-, -CHX<sup>a</sup>, -CX<sup>a</sup><sub>2</sub>-, -S-(CH<sub>2</sub>)<sub>m</sub>- and -N(R<sup>5</sup>)(CH<sub>2</sub>)<sub>m</sub>-, where m = 1-3, and X<sup>a</sup> is halogen.

The moiety ~~Ar~~ L<sup>1</sup> is preferably a 5-10 member aromatic structure containing 0-2 members of the group consisting of nitrogen, oxygen and sulfur which is unsubstituted or substituted by halogen up to per-halosubstitution and optionally substituted by Z<sub>n1</sub>, wherein n1 is 0 to 3.--



Please amend the second, third and fourth paragraphs on page 6 as follows:

D<sup>3</sup> --The aryl and heteroaryl moieties of B are more preferably of the formula:

Appl. No. 09/472,232  
 Amendment dated 4-03-03  
 Reply to Final dated 10/29/02

wherein  $\Psi$  M is selected from the group consisting of -O-, -S-, -CH<sub>2</sub>-, -SCH<sub>2</sub>-, -CH<sub>2</sub>S-, -CH(OH)-, -C(O)-, -CX<sup>a</sup><sub>2</sub>-, -CX<sup>a</sup>H-, -CH<sub>2</sub>O- and -OCH<sub>2</sub>- and X<sup>a</sup> is halogen.

D<sup>3</sup>



Q L is a six member aromatic structure containing 0-2 nitrogen, substituted or unsubstituted by halogen, up to per-halosubstitution and Q<sup>1</sup> L<sup>1</sup> is a mono- or bicyclic aromatic structure of 3 to 10 carbon atoms and 0-4 members of the group consisting of N, O and S, unsubstituted or unsubstituted by halogen up to per-halosubstitution. X, Z, n and n1 are as defined above, and s = 0 or 1.

In preferred embodiments, Q L is phenyl or pyridinyl, substituted or unsubstituted by halogen, up to per-halosubstitution and Q<sup>1</sup> L<sup>1</sup> is selected from the group consisting of phenyl, pyridinyl, naphthyl, pyrimidinyl, quinoline, isoquinoline, imidazole and benzothiazolyl, substituted or unsubstituted by halogen, up to per-halo substitution, or  $\Psi-Q^1-M-L^1$  is phthalimidinyl substituted or unsubstituted by halogen up to per-halo substitution. Z and X are preferably independently selected from the group consisting of -R<sup>6</sup>, -OR<sup>6</sup> and -NHR<sup>7</sup>, wherein R<sup>6</sup> is hydrogen, C<sub>1</sub>-C<sub>10</sub>-alkyl or C<sub>3</sub>-C<sub>10</sub>-cycloalkyl and R<sup>7</sup> is preferably selected from the group consisting of hydrogen, C<sub>3</sub>-C<sub>10</sub>-alkyl, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl and C<sub>6</sub>-C<sub>10</sub>-aryl, wherein R<sup>6</sup> and R<sup>7</sup> can be substituted by halogen or up to per-halosubstitution.--